

GUARANTEED ANALYSIS

| | 10-15-7 | |
|----------------------------------|----------|--------|
| Total Nitrogen | | 10.00% |
| 4.46% Ammoniacal 1 | Nitrogen | |
| 2.78% Nitrate Nitrog | en | |
| 2.76% Urea Nitrogen | í | |
| Available Phosphoric Acid (P2O5) | | 15.00% |
| Soluble Potash (K2O) | | 7.00% |
| Boron (B) | .002% | |
| Copper (Cu) | .05% | |
| Iron (Fe) | .10% | |
| Manganese (Mn) | .05% | |
| Zinc (Zn) | .05% | |

Derived From: Urea, Ammonium Nitrate, Ammonium Phosphate, Potassium Thiosulfate, Boric Acid, EDTA Copper, HEDTA Iron, EDTA Manganese and EDTA Zinc

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. In case of contact with eyes, flush immediately with copious amounts of water for 15 minutes. If irritation persists, consult a physician. Avoid contamination of skin and clothing. Wash with soap and water.

WARRANTY AND CONDITION OF SALE: Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. Buyer and user, therefore, acknowledges and assumes all risk and liability resulting from handling, storage, and use of this material.

AG BIO Logic, LLC Seneca, S.C. 29678

OVER THE TOP CORN

AUXILIARY NUTRIENT COMPOUND FOR CORN, SORGHUM, AND GRAIN

READ ENTIRE LABEL BEFOR USE. SHAKE WELL BEFORE USE.

DESCRIPTION

OVER THE TOP CORN is a proprietary formulation of auxiliary nutrients, (major and micronutrients), designed for foliar application on corn, sorghum, and other grains. The nutrient base is fortified with natural diterpines, extracts of ascophyllum nodosum, natural fermentation compounds, and enzymes.

COMPATIBILITY

OVER THE TOP CORN is tank mix compatible with a wide array of nutrient mixtures, pesticides, etc., but since local water conditions cannot be predicted a quart jar test should be conducted prior to tank mixing.

APPLICATION

Apply as a foliar mist to the foliage of the crop. Do not overspray to run off.

RATES AND DIRECTIONS

- <u>CORN and SORGHUM</u>: Apply 1.5-2.0 pints per acre, directed into the whorl at the 3-5 blade stage.
- BROADCAST GRAINS, (WHEAT, ETC.): Apply 1.5-2 pints per acre at jointing stage.

NET CONTENTS: 2.5 Gal. (9.45 Liters) 26.58 lbs / 12.08 Kg