

GUARANTEED ANALYSIS

3 - 2 - 3

m 137 00	
Total Nitrogen (N) 3.00	%
3.0% Ammoniacal Nitrogen	
Available Phosphoric Acid (P2O5) 2.00%	6
Soluble Potash (K2O) 3.00%	6
Boron (B)	

Derived From: Ammonium lactate. Potassium Pyrophosphates, Boric Acid, Copper, EDTA, Manganese EDTA, and Zinc EDTA

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. In case of contact with eyes, flush immediately with copious amounts of water for 15 minutes. If irritation persists, consult a physician. Avoid contamination of skin and clothing. Wash with soap and water.

WARRANTY AND CONDITION OF SALE: Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. Buyer and user, therefore, acknowledges and assumes all risk and liability resulting from handling, storage, and use of this material.

OVER THE TOP

FOR ADVANCED COTTON PRODUCTION READ ENTIRE LABEL FOR USE. SHAKE WELL BEFORE USE.

DESCRIPTION

OVER THE TOP is a proprietary formulation of major and micronutrients sequestered in a base of humic, fulvic, ulmic, and hymatheomelaic acids. It is fortified with extracts of Ascophyllum nodosum, Chondrus crispus, fermented plant extracts, enzymes, and amino acids. When used as directed as a foliar auxiliary program in your cotton production program, it can increase bloom and square initiation, boll set, and reduce fruit abortion due to environmental stresses.

COMPATIBILITY

OVER THE TOP should be tank mix compatible with most other nutrient compounds, insecticides, herbicides, etc. Since local water characteristics can vary greatly, a standard quart jar test is always recommended prior to tank mixing.

HOW TO APPLY

Apply with any equipment that delivers a fine even spray mist to crop canopy. Cover all foliage thoroughly. May be applied with ground equipment or aircraft. Best results will be obtained with spray volumes of 8-15 GPA with ground equipment and 2-3 GPA aerial.

RATE OF APPLCIATION:

Apply at any time after true leaves are formed.

- EARLY SEASON: To advance crop apply 1.5 pints (24 fl. oz.) per acre after true leaves are formed. Narrow band applications of 1.0 pint per acre are effective at this time.
- PIN HEAD SQUARE: Apply 1.5 pints (24 fl. oz.) per acre at early pin head square stage. Repeat in 12-21 days for heavy fruit set.
- MID BLOOM: Apply 1.5 pints (24 fl. oz.) per acre.

Any application will be beneficial. At least (2) applications are recommended.

AGBIO Logic, LLC Seneca, S.C. 2968 NET CONTENTS: 2.5 Gal (9.45 liters) NET WEIGHT: 21.65 lbs (9.84 Kg)